




## TECHNIQUE OF PERIODONTAL PROBING


- Modified pen grasp
- Walked around the teeth parallel with the root surface
- Measure at 6 points of each tooth
- Light probing force (25g)

## WHICH FACTORS DO INFLUENCE THE PROBING ?










	120	10
	PERIODONTA	L ASSESSMENT
1112	MARIAN CLARMEN	♦ FMPS: <b>51 %</b>
adatada ventrada	Deren har an harde	utile coprestic
		⊗FMBS: <b>62 %</b>
1	CONTRACTOR AND A CONTRACT	⊗ PPD≥5mm: <b>46 %</b>
The provide state of the second state of the s		and mr. Dr. H


DIA	GI	N	ő	ST	S	.1	PT			31	T	0	SI	S			
				2011													
♦ Diagnos:	is: Pe	erio	don	titis	ch:	roni	ca										
♦ Prognosi																	Aug
hopeless	Τ							50	12								
doubtful	x	x	x	x		.0	¢.`						x	x	x	x	
stable					x	x	x	x	x	x	x	x					28
	8	7	6 (	5	4	3	2	1	1	2	3	4	5	6	7	8	
stable			x	x	x	x	x	x	x	x	x	x	x				
doubtful	x	x							30	~					x	x	

antil	
our c	OF COMPREHENSIVE PERIODONTAL THERAPY
	CPITN index Systemic / Ac.sta Treatment of MH and acut lesions (e.g. abscess, NUG/NUP)
	Full perio chart: PPD, REC, CAL, FMPS, FMBS, furcation (I-III), mob (1-3)
<b>5.</b> 6. 7. 8.	Case presentation, consequences of no treatment Oral hygiene instructions, smoking cessation counseling Root surface debridement, elimination of plaque retentive factors, temporary splinting, fluoride application, monitoring/improving OH Re-assessment at 6-8 weeks (full perio chart), Corrective Tx plan
	Periodontal surgeries (PPD≿5mm) Re-assessment (perio chart PPD should be ≤4mm) Definitive prosthetic, implant, ortho Tx
	Periodontal supportive care (risk analysis)














B







# TREATMENT PHASES

#### OF COMPREHENSIVE PERIODONTAL THERAPY

#### CPITN index

Treatment of MH and acut lesions (e.g. abscess, NUG/NUP)

Full perio chart: PPD, REC, CAL, FMPS, FMBS, furcation (I-III), mob (1-3) Tooth by tooth prognosis (secure, doubtful, hopeless)

- Case presentation, consequences of no treatment
- . Oral hygiene instructions, smoking cessation counseling
- Root surface debridement, elimination of plaque retentive factors, temporary splinting, fluoride application, monitoring/improving OH
- Re-assessment at 6-8 weeks (full perio chart), Corrective Tx plan

### Periodontal surgeries (PPD≥5mm)

- Re-assessment (perio chart PPD should be Definitive prosthetic, implant, ortho Tx
- 2. Periodontal supportive care (risk analysis)<sup>Supportiv</sup>


# PERIODONTAL SUPPORTIVE CARE

### Concept of recall:

- Succesfull complex periodontal treatment
- Adequate individual motivation and oral hygiene
- Regular professional mechanical cleaning in *every 2-12* months depending on the clinical case, risk factors, individual oral hygiene and manuality

### Motivation and instruation every time!

9



- Uses electric brush only
- Occasionally mouthrinses



