

Dental attendance of children suffering from general disease

Dr. Beck Anita, Dr. Bálint Réka, Dr. Macsali Réka

Semmelweis University,

Department of Paediatric Dentistry and Orthodontics




SEMMELWEIS
UNIVERSITY 1769

Dental attendance of children

- Diseases of teeth and oral mucosa in (otherwise) healthy children
- (Otherwise) healthy children, met traumatic injuries
- Children suffering from infectious diseases
- Syndromes and other congenital or genetic disorders in children
- Children suffering from mental illnesses
- **Children suffering from general systematic diseases**

Dental attendance of children suffering from general systematic disease

- Diabetes Mellitus
- Kidney
- Cardiovascular
- Haematological
- Respiratory, asthma
- Allergy
- Neurological




At present, diabetes cannot be cured, but it can be treated effectively and can be asymptomatic if blood glucose levels are close to normal.

- Diet
- Exercise
- Medicine

News-Medical

Diabetes Mellitus I.

- IDD (I. Typ)
 - 10%
 - Manifestation under 30 („juvenile DB”)
 - Body: normal-astheniac
 - Digestion: labile
 - ID



Diabetes Mellitus II.

- NIDD (II. Typ)
 - 90%
 - Manifestation over 40 („adult DM”)
 - Body: adiposus
 - Digestion: relativ stabil
 - NID

Diabetes Mellitus

- Oral manifestation:
 - Saliva glucose level increase
 - Caries predisposition
 - Xerostomy
 - Burned feeling of mucosa
 - Oral ulceration
 - Paro disease
 - Paro abscess
 - Candidiasis, infection

Potential problems at the dental praxis

- Not treated DB: upset of metabolism (pl. Stress, workout, starving, insulin) => hypoglycaemia, ataxia, tachycardia, hipertensio, fear of death, strange speech, acetonic breath (looks drunk)
- Infection predisposition (Str. A Haemol., Staph)
- Woundhealing difficulties
- Cardiovascular (hypertonia, cardiac insufficiency, angina p., infarctus)
- Optic (retinopathia, blindness)
- Nephrological (renal failure)
- Neurological (periferial neuropathy)

Dental treatment

- Consultation with general doctor/ specialist!!!
- Aim: PREVENTION
- Need of steroid?
- Need of AB profilaxis?
- Don't use anesthetic or retraction thread with tonogen
- Don't use medicaments containing sugar, sweetener
- Drug interactions: NSAID, acetyl, tetracyclin, asparagin
- Avoidens of hypoglycaemy: eat before (little), take insulin before if needed, morning treatment

Kidney diseases

- Nephrosis syndrome (Proteinuria)
- Dental developmental and eruption disorder
- Drug metabolism change depending on renal function:
 - NSAID
 - Benzodiazepines
 - Steroids
- Aim: PREVENTION

Cardiovascular diseases

- Often part of a syndrome
- Increased bleeding predisposition
- Increased infectious predisposition
- Often hospitalised
- Deciduous teeth: caries
- Permanent teeth: hypoplasia

Pinterest

Congenital Cardiac diseases

- 15% sy (Down 70%)
- Predisposition: rubella, diabetes, alcoholism, drugs, smoking)
- Cyanotic:
 - Early surgery
 - Venous blood (from the left to the right)
 - Fallot-tetralogy
 - Pulmonary atresia
 - Vascular transposition
- Acyanotic:
 - Obstructive: aortic stenosis, pulmonary stenosis, coarctation of aorta
 - Shunt: atrial, ventricular, atrioventricular septum defect, Pers. Ductus Botalli

Cardiac diseases after birth

- Rheumatic fever
- Myocardium, pericardium inflammations
- Arrhythmies
 - Additional impulse conduction ways => AV nodus=> early stimulus=> tachycardia, rhythm disorders
- Other

Cardiavascular diseases

Dental treatment

- Consultation
- Aim: PREVENTION (diet, oral hygiene, F⁻, FS), controll
- Whole mouth sanation before heart sugery
- Need of AB profilaxis? Inform parents about IE
- Every invasive intervention should be done under profilactic defence in high risk patients, otherwise AB not indicated
- Drug type, form, dose, repeat
- Before invasive intervention CHX

IE High risk patients

- After correction of cardiovascular defect, if there was an implantation (valve replacement, cardiac catheter)
- IE in anamnesis
- Mitral prolapsus with regurgitation
- Severe congenital developmental cardiac disorder (cyanotic or regurgitational)

AB-prophylaxis: 20mg/bwkg Penicillin

50 mg/bwkg clindamycin

15 mg/bwkg macrolid

High risk patients

Dental treatment

- cofferdam, matrice, wedge, sulcusexpander
- Paro probe, surgery, depuration
- Rootcanal treatment
- Extr, repl, impl, sep.
- Intraligamentar injection (Tonogen not contraindicated)
- Not dangerous:
 - Pulpotomy (?)
 - Abscess incision
 - Removal of suture

Haematological diseases

- Coagulation disorders
- Quantitative disorders of red blood cells
- Quantitative disorders of white blood cells

Pinterest

Coagulation disorders

- Genetic
 - Haemophilia A
 - Haemophilia B
 - Von Willebrand sy (VIII. factor missing, thrombocyta-adhesion decrease)
- Medicine side effect
- Hepatic diseases
- Infections reported commonly
- Other severe diseases (f.e: leukaemia)

Thrombocyta count

- Norm: $300 \times 10^9/l$, minimally acceptable $50 \times 10^9/l$
- Thrompocytopaenia:
 - Producing by bone marrow decreases (drug, aplast anaemia)
 - Idiopathic
 - Virus or rubella vaccination IgG (4-6 weeks spontan healing, skin- or mucosa petechia purpura)
- 75 g/l wound bleeding
- 25 g/l spontan bleeding
- 15 g/l skin petechia
- 5 g/l oral petechia

Protrombin rate

- For Protrombin-Trombin transformation
- Measurement: Quick - time (tromboplastin time)
 - Norm: 100% (80-120%)
 - Minimally acceptable: 40% (INR=2,5)
INR (International Normal Relation)
 - Healthy Quick time (100%) proportional to the measured Quick
 - Quick decreases, INR increases
 - Pl. If PR= 50 %, INR=2
 - Maximally acceptable: 2,5 (Quick: 40 %)

Coagulation disorders

Dental treatment

- Consultation
- Aim: PREVENTion (diet, oral hygiene, F-, FS), control
- Whole mouth rehabilitation
- Stand clear of invasive intervention, if there is high risk of bleeding: laboratory blood test factors of coagulation and thrombocyte rate (TCT)
- Drug type, form, dose, repeat

Coagulation disorders

Dental treatment

Platelet count or function decreases => no coagulum

- => bleeding after trauma (haemophylia it lasts for 4h)

- => prolonged bleeding episodes, spontan gum bleeding

- => predisposition to infections

No Szokoloczy! (maybe after factor supplementation)

No extraction, rather endodontic treatment (Except deciduous teeth)

In case of postoperative bleeding

- Preoperatively: provide with written and verbal information, what to do if bleeding doesn't stop in 1-2 hours
- Diet, oral health, No NSAID No!
- Local pressure
- Suture
- Spongostan with trombin
- Oxycellulose, Collagen
- Drug: Tranexamic acid (Exacyl tabl., rinsing-antifibrinolytic)
- Transfusion

Changes in the amount of red and white blood Cells

Red blood Cells

- Anaemia
- Polycythaemia

White blood Cells

- Leukopaenia
- Leukocytosis
- Leukaemia (80% ALL, CLL, 20% AML, 1% CML)
- Lymphoma (HL,NHL)

Anaemia

- Erythropoiesis dysfunction: - lack of Fe²⁺, B12-vit, C-vit., folic acid
- bone marrow functional disorder (aplast., AI, malign, burn)
- Increased death of red blood Cells: - extrinsic: transfusio, drug
- intrinsic: ensimopathia, AI, thalassemia)
- Blood loss: acut/chronic bleeding
- **Symptoms** Glossitis (papilla atrophica), tongue burn, pain, stomatitis, cheilitis angularis
- Forms: normocyta, microcyta, macrocyta
normochrom, hypochrom, hyperchrom

Leukaemia

- Chronic: normal leukocyte count increases
- Acute: blast cells pile up
- Symptoms: pale-faced, anorexia, hepatosplenomegalia, lymphadenopathy, infections, anaemia
- **Oral symptoms** bleeding, petechia, ANUG, sore throat, swollen glands, xerostomia, foetor => „leukaemic mouth”
- Parodontal defect, bone deformation, tooth moves
- Aplasia, microdontia, radicleless teeth

Leukaemia

Dental treatment

- Only in remission!
- Consultation with general doctor/ specialist!!!
- Aim: PREVENTION diet, oral hygiene, F-, FS), controll
- Fungal, viral, AB profilaxis (Nystatin, Acyclovir, CHX)
- Salive supplement: Biotene
- Whole mouth sanitation
- Get clear of invasive intervention, no Szokoloczy
- Medicine type, form, dose, repeat

Changes in the amount of red and white blood cells, Dental treatment

- High caries risk
- Consultation
- Aim: PREVENTION (diet, oral hygiene, F-, FS), controll
- Whole mouth sanitation
- Stand clear of invasive intervention without hospital background, if there is high risk of complication: laboratory blood test, profilaxis
- Drug type, form, dose, repeat

Changes in the amount of red and white blood cells, Dentistry connected complication

- Immundeficiency
- Infection predisposition, mucositis
- Ulceration
- Slow wound healing
- Coagulation disorders
- Pain, bleeding => leave oral hygiene routine
- Drugs: nausea, neuropathic symptoms

Respiratory diseases

- Asthma
 - Stress provoking factors
 - Preoperative 1-2 puff inhalator (decreases saliva, lactose!)
 - Ventolyn corticosteroid-powder
- Allergy, tonsillitis, nose polip, septumdefect
- Cystic Fibrosis (AR)
 - Exocrin glands multisystematic disease ,hepatic cirrhosis, haemophilia
 - Frequent eating => cariesrisk
 - Not allowed to treat in general anesthesia, AB profilaxis needed (tetracyclin)

Epilepsy

- Drug hydantoin => gingivahyperplasia, xerostomia
=> drowsiness, coordination decreases
- Prevention of epileptic seizure: bensodiasepine
- In case of epileptic seizure: Seduxen (im. 10mg , Rectal)
- Preparation of assistency, trigger minimalisation
- Don't provide them with removable appliances
- Total luxation/ avulsion no replantation
- Fix nonceramic prosthetic devices

