




*250 years of EXCELLENCE
in medical education,
research & innovation
and healthcare*

Pathology of Endocrine Organs

Dr. Kovács Kristóf Attila

29th March 2021



Pathology of Endocrine Organs Part II.

Outline of the lecture

1. Basics of endocrine anatomy and pathophysiology (=hormonal dysfunctions)
2. Pathology of endocrine organs
 - Pituitary gland
 - Pineal gland
 - *Thyroid gland (see next lecture!)*
 - Parathyroid glands
 - Adrenal glands
 - Endocrine pancreas
 - Sex-cord stroma
3. General aspects of neuroendocrine tumors of non-endocrine organs (DNES)
4. Syndromes associated with endocrine tumors


Anatomy

Endocrine tissues

Neuroendocrine tissues

Origin

- *Endocrine: thyroid follicles, adrenal cortex, sex-cord*
- *Neuroendocrine (tumors with neurosecretory features): pituitary gland, thyroid C-cells, parathyroid glands, adrenal medulla, gastro-entero-pancreatic and other neuroendocrine cells of respiratory, urinary tract, skin etc. (GEP/DNES)*




Pathophysiology of endocrine organs

Endocrinological classification (pathophysiology)


- Causes and consequences of decreased/no hormone function (=hypo...ism)
 - Underproduction
 - No effect
- Causes and consequences of increased hormone function (=hyper...ism)
 - Overproduction

Regulation of endocrine organs

- With a regulating hormone



- Without a regulating hormone



Special definitions in endocrine pathology


- **Congenital disorders=enzimopathies**

- Inappropriate enzyme production
- No effective enzyme function

- No feedback

- RESULT: Hyperplastic endocrine organ with hypo...ism


- Examples: dyshormonogenetic goiter, adrenogenital syndrome



Special definitions in endocrine pathology

- **Adaptive disorders**

- Primary hyperfunction = autonomous hormone production
- Secondary hyperfunction = excessive release of regulating hormon/factor
 - Central
 - Ectopic (paraneoplasia)
- Tertiary hyperfunction = secondary hyperfunction with an autonomous population



Pathology of endocrine organs

Morphological classification (pathology)

- Congenital disorders
- Adaptive disorders
- Circulatory disorders
- Inflammatory disorders
- Neoplasia

General aspects in endocrine pathology

- **Circulatory disorders**

- Very rare= small organs with rich vascularisation
- Ischemia can occur in the pituitary gland due to its limited space

- **Inflammations**

- Almost always of autoimmune etiology
- T-cell mediated (type IV) eg. diabetes mellitus type 1, autoimmune adrenalitis
- Antibody mediated (type II) → acts as a regulating hormone eg. Graves disease
- Infections affecting endocrine organs are very rare

General aspects in endocrine pathology

- **Neoplasias**


Origin

- Endocrine: thyroid follicles, adrenal cortex, sex-cord
- Neuroendocrine (tumors with neurosecretory features): pituitary gland, thyroid C-cells, parathyroid glands, adrenal medulla, gastro-entero-pancreatic and other neuroendocrine cells of respiratory, urinary tract, skin etc. (GEP/DNES)


Dignity

- Classic cytomorphologic criteria of malignancy is NOT valid in endocrine organs!
- Atypia, polymorphism, bizarre cells is more often found in hyperactivity than malignancies!
- *Exemption: papillary thyroid carcinoma* → diagnosis based on special cytological features
- Definition of endocrine malignancy in general: **local/vascular invasion+metastasis**
- Most neuroendocrine tumors are potentially malignant irrelevant to their microscopic morphology (see DNES tumor classification)


1 Adrenal cortical adenoma with bizarre cells




2 Hashimoto thyroiditis with Hürtle cells




3 Parathyroid adenoma with bizarre cells



4 Medullary thyroid carcinoma




Pituitary gland *(p1098)*




- Anatomy
 - Anterior lobe (adenohypophysis)
 - Posterior lobe (neurohypophysis)
- Note: local mass effect may cause visual field abnormalities
- Function:
 - Anterior lobe
 - Somatotroph cells: GH
 - Lactotroph cells: prolactin
 - Corticotroph cells: ACTH
 - Thyrotroph cells: TSH
 - Gonadotroph cells: FSH, LH
 - Posterior lobe: ADH, oxytocin

Hyperpituitarism


- >99% adenomas
 - Microadenoma: <1 cm, macroadenoma: >1cm
 - Classification based on hormon secretion
 - Most common: prolactinoma galactorrhea
 - Somatotroph adenoma gigantism, acromegaly
 - Corticotroph adenoma central Cushing syndrome
 - Other rare variants (gonadotroph, thyrotroph, nonfunctioning)



Morphology



Macro
Adenoma





Micro

Normal: mixture of:

- acidophils (GH&PRL)
- basophils (ACTH, TSH, FSH&LH)
- Chromophobes


Adenomas: monotonous population



Hypopituitarism

- Physical damage
 - Mass lesion: adenoma
 - Trauma
 - Surgery, irradiation
- Vascular disorders
 - Pituitary apoplexy (rarely sudden death)
 - Sheehan syndrome (ischemic necrosis in pregnant women)
- Genetic defects
- Hypothalamic hormone deficiency (eg. suprasellar tumors)
- Infections (rare, granulomatous, eg. TB, sarcoidosis)

Any sort of pituitary damage may result empty sella syndrome




Posterior pituitary syndromes

- Hyperfunction
 - Syndrome of inappropriate ADH (SIADH)
 - Generally ectopic ADH secretion=paraneoplasia (small cell lung carcinoma)
- Hypofunction
 - Diabetes insipidus
 - Trauma, tumor etc.

Pineal gland


- Tumors



- Pinealoma – deriving from pinealocytes, no special hormonal syndrome
- Interestingly most of pineal gland tumor are of germ cell origin (eg. teratoma)

Parathyroid glands (*p1126*)

- Anatomy
 - 2 x paired organs
 - Rarely mediastinal location (intrathyroidic)
 - Histologically 3 different cell types (chief, water clear, oxyphil)
- Function
 - Parathormone release
 - Directly controlled by blood Ca^{2+}
 - Very short half life – Ca^{2+} level is able to monitorize PTH release





Hyperparathyroidism, hypercalcemia


- Primary (PHPT)
 - 90% adenoma, 9% hyperplasia (MEN), 1% carcinoma
 - Usually solitary lesion
 - Preoperative diagnosis: Radionuclide scan/FNAB
 - Surgery: intraoperative Ca 2+ test!
- Secondary
 - Cause: excessive Ca 2+ loss
 - Chronic renal failure
 - Hyperplastic change in all glands
- Malignancy associated hypercalcemia – PTH related protein (PTH-rP)
 - Common paraneoplasia with poor prognosis
 - Solid tumors: lung-head&neck squamous cell carcinoma, breast carcinoma, renal cell carcinoma
 - Hematologic malignancy: multiple myeloma

Morphology

normal parathyroid gland




adenoma




hyperplasia

Micro morphology




normal



adenoma

Consequences of hyperparathyreoidism

- Increased osteoclast activity
 - Skeletal lesions: osteitis fibrosa cystica, „brown tumor”
- Elevated Ca^{2+} level
 - Repeating nephrolithiasis
 - Gastrointestinal malfunction (constipation)
 - CNS: depression
 - Neuromuscular: weakness
 - Cardial disorders: valvular calcification



Hypoparathyroidism

- Generally iatrogenic (total thyroidectomy)
 - Autoimmune parathyroiditis
 - Genetic disorders
 - Aplasia
-
- Consequences: decreased Ca^{2+} level
 - Tetany, anxiety-depression, ocular and dental disorders, heart conduction abnormalities etc.



Adrenal glands (p1148)

- Anatomy

- Paired organs
- Rarely ectopic=eg. funicular
- Different development (mesodermic&neuroectodermic)

- Function

- Cortex: steroid hormones
 - Glucocorticoids
 - Mineralocorticoids
 - Sex steroids
- Medulla: catecholamins



Adrenocortical hyperfunction

- **Cushing syndrome = hypercortisolism**

- Central (=Cushing disease)
 - Pituitary corticotroph adenoma
 - Secondary hypercortisolism (ACTH↑, cortisol↑)
 - Diffuse adrenal hyperplasia
- Peripheral Cushing syndrome
 - Primary adrenal nodular hyperplasia or adenoma, rarely carcinoma
 - Primary hypercortisolism (ACTH↓, cortisol↑)
 - Adrenal cortex atrophy
- Ectopic Cushing syndrome
 - Paraneoplasia, eg. SCLC
 - Secondary hypercortisolism (ACTH ↑ ↑ ↑, cortisol↑)
 - Diffuse adrenal hyperplasia
- Exogenous Cushing syndrome
 - Glucocorticoid medication generally for immunosuppressive purpose

- **Conn syndrome = hyperaldosteronism**

- primary adrenal adenoma
- rarely nodular hyperplasia
- Secondary hyperaldosteronism
 - Increased renin level: renal hypoperfusion, pregnancy
 - Renin producing tumors: very rare

- **Hyperfunction of sex hormones**

- Virilization (masculinization)
 - amenorrhea, infertility, male distribution of hair=hirsutism, voice changes, clitoris hypertrophy
- Feminization
 - Loss of hair, decreased libido etc.

Cushing's Disease or Syndrome Symptoms



Hyperandrogenism




Tall stature

Slightly feminized physique

**Mildly impaired IQ
(15 points less than average)**

Tendency to lose chest hairs

Female-type pubic hair pattern



Frontal baldness absent

Poor beard growth

**Breast development
(in 30% of cases)**

Osteoporosis

Small testes

Morphology of adrenal cortical lesions



adenoma

primary=nodular





hyperplasia

secondary=diffuse



Microscopic morphology



Adrenocortical insufficiency

Primary

- Genetic abnormality
 - adrenogenital syndrome=21 hydroxylase deficiency
 - mineralocorticoids ↓, glucocorticoids ↓, sex steroids ↑
 - ACTH ↑ ↑ ↑
 - Super-sized adrenals
- Congenital hypoplasia
- Acute hemorrhagic necrosis
 - =Waterhouse-Friderichsen syndrome (meningococcus sepsis)
- Primary chronic adrenocortical insufficiency= Addison disease
 - Autoimmune adrenalitis
 - Infections (fungal, TB)
 - Metastatic cancer, lymphoma

Secondary

- ACTH deficiency
 - Hypopituitarism
 - Long term steroid administration=atrophy






Figure 34
(Figures 34 and 35 from same patient)
CONGENITAL ADRENAL HYPERPLASIA
Gross photograph of kidney and adrenal from a patient with congenital adrenal hyperplasia who died at the age of 6 weeks. Combined adrenal weight was 22 g. 17-ketosteroid urinary excretion in 24 hours was increased to greater than 10 times normal. This is probably an example of severe 21-hydroxylase defect. Actual size. (Courtesy of the Department of Pathology, University of Leeds, Leeds, England.)


Waterhouse -
Friderichsen
syndrome



Adrenal medulla

- Pheochromocytoma
 - >90% hormonally active=secondary hypertension
 - >90% benign clinical course (no evident histological signs of malignancy)
 - Frequent germline genetic abnormalities (see later)



Morphology



Ganglion-like chief cells and sustentacular cells form „zellballen“

Embryonal type medullary tumors

- Neuroblastoma (see childhood tumors)
- Matured form: ganglioneurinoma – incidental, hormonally inactive tumor in adults




Hormonally inactive adrenal tumors

- Incidental cortical adenoma
 - Most frequent autopsy finding within the adrenals
- Metastatic malignancies
 - Lung-, kidney-, breast-, colorectal carcinoma, melanoma



Endocrine pancreas

- Anatomy
 - Islets of Langerhans are not visible macroscopically
- Function
 - β -cells: insulin production
 - α -cells: glucagon production
 - δ -cells: somatostatin
 - Other hormone products: pancreatic polypeptide, vasoactive intestinal polypeptide, gastrin



Hyperfunction of endocrine pancreas

- Always primary – generally hormonally active neuroendocrine tumors
 - Hyperinsulinism: insulinoma (generally benign), rarely islet hyperplasia (nesidioblastosis) → hypoglycemia
 - Glucagonoma → diabetes, characteristic skin lesion (necrolytic migratory erythema)
 - Gastrinoma → Zollinger-Ellison syndrome
 - VIPoma → WDHA syndrome = watery diarrhea, hypokalemia, achlorhydria)
- General tumor features: see NET classification

Hypofunction of endocrine pancreas

- Diabetes mellitus (DM) is a group of metabolic disorders caused by hyperglycemia which is a result of defect in insulin secretion or insulin action or both
- Hyperglycemia is not a morphological disorder thus DM is a clinical diagnosis
- Classification of DM also based on clinical tests – morphology of endocrine pancreas is not examined routinely
- Pathology studies focus on the late morphological consequences of DM (generally faced them in the autopsy hall)

Classification of DM (p1132)

- Mixed insulin resistance and β -cell dysfunction: type 2 DM
- Inflammatory
 - Autoimmune T-cell mediated (hypersensitivity type IV): DM type 1
 - Infective: very rare (CMV)
- Exocrine pancreas defect (pancreatogenic DM)
 - Chronic pancreatitis
 - Total pancreatectomy
- Gestational DM
- Endocrinopathies
- Drugs
- Genetic causes

Pathogenesis of complications in DM (p1138)

Basic problems resulted from


1. Glycation
2. PKC activation and
3. Intracellular hyperglycemia

The problem comes from the formers → degeneration of:

1. Arterial intima
 2. Arterioles
 3. Basement membrane
 4. Retina
 5. Nerves
- + Increased risk of infections


Target organs affected by DM and its morphology

(p1139-43)





Macrovascular Disease

- Accelerated atherosclerosis
 - DM is one (but not the only!) of the most important risk factors of atherosclerosis
- PAD: Peripheral Artery Disease
 - Commonly affect legs



Cardial complications

- AMI – accelerates coronary sclerosis
- Valvular and mitral annulus calcification



„Diabetic foot”





- Gangrene and skin ulcers resulted by
 - PAD
 - Microangiopathy
 - Neuropathy
 - Infections




Renal complication

- Micro- and macroangiopathy
- Glomerular damage
 - Nodular glomerulosclerosis = Kimmelstiel-Wilson disease
 - Diffuse mesangial sclerosis
- Pyelonephritis
 - Acute PN with necrotizing papillitis
 - Chronic PN
- End stage kidney

nephrosclerosis




Kimmselstiel-Wilson disease (see nephropathology lecture)



Eye disorders

- Diabetic retinopathy
 - Proliferative microangiopathy
 - Microhemorrhages
 - Fibrosis
- Cataracta
- Glaucoma



Infections in DM

Rhinocerebral mucormycosis



TABLE 1. CLINICAL FEATURES, DIAGNOSIS, AND CAUSATIVE ORGANISMS OF SELECTED INFECTIONS IN PATIENTS WITH DIABETES.

INFECTION	CLINICAL FEATURES	DIAGNOSTIC PROCEDURE*	ORGANISMS	COMMENTS
Respiratory tract Community-acquired pneumonia	Cough, fever	Chest radiography	<i>Streptococcus pneumoniae</i> , <i>Staphylococcus aureus</i> , <i>Haemophilus influenzae</i> , other gram-negative bacilli, atypical pathogens	Pneumococcal infection carries a higher risk of death in diabetic than in nondiabetic patients
Urinary tract Acute bacterial cystitis	Increased urinary frequency, dysuria, suprapubic pain	Urine culture	<i>Escherichia coli</i> , proteus species	Bacteriuria more common in diabetic than in nondiabetic women
Acute pyelonephritis	Fever, flank pain	Urine culture	<i>E. coli</i> , proteus species	Emphysematous infection should be considered
Emphysematous pyelonephritis	Fever, flank pain, poor response to antibiotics	Radiography or CT scanning	<i>E. coli</i> , other gram-negative bacilli	Emergency nephrectomy often required
Perinephric abscess	Fever, flank pain, poor response to antibiotics	Ultrasonography or CT scanning	<i>E. coli</i> , other gram-negative bacilli	Surgical drainage usually required
Fungal cystitis	Same as for acute bacterial cystitis	Urine culture	Candida species	Difficult to distinguish colonization from infection
Soft tissue† Necrotizing fasciitis	Local pain, redness, crepitus, bullous skin lesions	Radiography or CT scanning	Gram-negative bacilli, anaerobes (type I), or group A streptococci (type II)	High mortality; emergency surgery required
Other Invasive otitis externa	Ear pain, otorrhea, hearing loss, cellulitis	Clinical examination, magnetic resonance imaging	<i>Pseudomonas aeruginosa</i>	Prompt otolaryngologic consultation recommended
Rhinocerebral mucormycosis	Facial or ocular pain, fever, lethargy, black nasal eschar	Clinical examination, magnetic resonance imaging, pathological findings	Mucor and rhizopus species	Strong association with ketoacidosis; emergency surgery required
Abdomen Emphysematous cholecystitis	Fever, right-upper-quadrant abdominal pain, systemic toxicity	Radiography	Gram-negative bacilli, anaerobes	High mortality; gallstones in 50%; emergency cholecystectomy required

*CT denotes computed tomography.

†Foot infections are described in detail in Table 3.

Sex cord neoplasias (*p1050 and 992*)

Hyperfunction of sexual hormones: see adrenal cortical hyperfunction

- Ovarian tumors

- Granulosa-Theca cell tumors
 - Hyperestrogenism may cause endometrial HP and carcinoma!
- Fibroma/thecoma
 - Usually incidental, hormonally inactive
- Sertoli-Leydig cell tumors
 - Hyperandrogenism



- Testicular tumors

- Leydig cell tumor
 - Precocity in childhood
- Sertoli cell tumor

Common macroscopic feature:
yellowish color




General aspects of neuroendocrine tumors of non-endocrine organs (DNES)

- Deriving from dispersed neuroendocrine cells of epithelial tissues (aka APUD cells)
- Development:
 - Foregut: thymus, trachea, bronchus
 - Midgut: stomach, small intestine, appendix, pancreas, biliary tract
 - Hindgut: colon, rectum
- Diagnostic criterion: presence at least 1 NE marker immunohistochemically
- Gastro-Entero-Pancreatic NeuroEndocrine Tumors (GEP-NET)
 - WHO classification of GEP-NET 2010
 - Neuroendocrine tumor, grade 1-3 (based on mitotic count/Ki67 index)
 - Neuroendocrine carcinoma grade 3 (based on morphology)
- Neuroendocrine tumors of the lung
 - WHO classification of lung tumors 2015
 - Typical/atypical carcinoid
 - Small cell lung carcinoma (SCLC)
 - Large cell neuroendocrine carcinoma
- Others
 - Urinary tract, skin, breast etc.

Hormonal activity in neuroendocrine tumors

- Stomach/duodenum: gastrin (ZES)
- Pancreas: insulin, glucagon, gastrin (ZES), somatostatin, VIP
- Small bowel: **serotonin**
- Lung: **serotonin**, ACTH (Cushing), PTH



Survival of GEP-NETs

- Much better than conventional carcinomas
- Long survival even with liver mets


SEER Stage	5-Year Relative Survival Rate
Localized	98%
Regional	93%
Distant	67%
All SEER stages combined	94%

Macroscopic morphology of neuroendocrine tumors

- Relatively small size
- Yellowish color
- Very tough consistency




Small intestine



Bronchus



Appendix




Pancreas

Microscopic morphology of neuroendocrine tumors


Neuroendocrine tumors:

1. Insular structures common
2. Salt&pepper chromatin
3. Plasmocytoid appearance
4. Low mitotic count (in G1-2)
5. No necrosis



Poorly differentiated neuroendocrine carcinoma (small cell type)


1. Hyperchromasia
2. Polymorphism
3. High mitotic count
4. Necrosis




Grading of GEP-NETs

- Grade 1: Ki 67: <2%, mitoses: <2/10 HPF
- Grade 2: Ki 67: 2-10%, mitoses 2-10/10 HPF
- Grade 3: Ki 67: >20%, mitoses >10/10 HPF

Presence of neurosecretory vacuoles



IHC



EM

Overview of NE neoplasias in different organs

Organ	<u>Neuroendocrine tumors</u> Clinical behaviour: slow progression even in metastatic stage	<u>Neuroendocrine carcinoma</u> Clinical behaviour: poor prognosis, rapid progression
Pancreas	NET grade 1,2 Insulinoma: practically benign	NEC grade 3: poor prognosis
Stomach, duodenum	NET grade 1,2 - MEN 1 & atrophic gastritis associated: good prognosis - Sporadic: worse prognosis	
Small bowel	NET grade 1,2,3: high metastatic potential	
Colon	Rectum: grade 1,2: good prognosis	Proximal colon: NEC grade 3: poor prognosis
Appendix	NET grade 1: usually incidental finding in appendectomy: good prognosis	
Lung	Typical/atypical carcinoid: indolent clinical course	NEC (small/large cell): poor prognosis Frequent paraneoplastic hormon secretion
Urinary tract		NEC: poor prognosis
Skin		Merkel cell carcinoma: highly aggressive
Breast	Generally ER positive Prognosis not differ from usual breast cancer	

Syndromes associated with endocrine tumors

(p1161)

Syndrome	Mutated gene	Tumor types	Prognosis
Multiple endocrine neoplasia (MEN) type 1	MEN1	„3Ps“ Parathyroid (PHPT) Pancreas Pituitary Gastric and duodenal gastrinomas (multiple)	Poor if NEC occur (mainly of pancreas)
MEN type 2	RET	2A: Parathyroid hyperplasia, pheochromocytoma, medullary thyroid carcinoma - Preventive thyroidectomy needed in childhood - Pheochromocytomas are usually benign 2B: plus neuromas and marfanoid habitus Familial medullary thyroid cancer - Only MTC occur	Good if preventive thyroidectomy performed
VHL	VHL	Phaeochromocytoma	Other cancer types (renal cell carcinoma)
Neurofibromatosis	NF1	Phaeochromocytoma	
Familial paraganglioma (more types)	SDH (succinil dehydrogenase)	Phaeochromocytoma	Frequently malignant phaeochromocytoma

Summary

- Syndromes associated with hyper and hypofunction of endocrine/neuroendocrine organs
- Paraneoplastic syndromes
- Specialities of tumors in endocrine/neuroendocrine system

attila.drkovacs@gmail.com